

Faunistic spectrum of terraneous mollusks of Uzbekistan and bordering territories

Feruza Gaibnazarova^{1*}

¹Gulistan State University, Republic of Uzbekistan

Abstract. The article presents a list of species of 11 families, their ecological and geographical characteristics. The features of the distribution of terrestrial mollusks in connection with the vertical geobotanical belt in the conditions of the mountain systems of Uzbekistan and adjacent territories have been studied. The ecological groups of terrestrial molluscs have been determined in relation to the moisture factor and habitats. The faunistic spectrum of terrestrial mollusks was studied based on the materials of our collections, the material for this work was collected in Karzhantau, Ugam, Pskem, Chatkal, Talass, Kuramin, Fergana, Alai, Turkestan, Zarafshan, Gissar, Nurata, Kugitagtangtau in Chirchiko-Akhangaran, Zarafshan, Kashkadarya and Surkhan-Sherabad valleys, as well as extensive collections of the Zoological Institute of the Russian Academy of Sciences, the Zoological Museum of Moscow University

1 Introduction

The malacofauna of Central Asia is very diverse. It is distinguished by its rich endemism of species and generic rank, which is associated with its southern position and the diversity of the natural ecological environment. Currently, there are approximately more than 200 species of terraneous mollusks known in the fauna of Central Asia, whose systematic position has been studied quite well. At the same time, many aspects of their lives: biology, ecology, distribution, economic significance are investigated fragmentally. It should be noted that this aspect is affected in one way or another in many works devoted to the research of the Central Asian malacofauna. As a result, data on the species composition of terraneous mollusks were significantly replenished due to the inclusion of a number of species, new to the fauna of Central Asia; specified the boundaries of the ranges of a number of species; revision of the taxonomic position of certain forms; identified the identity of malacofaunas of different high-altitude zones. It should be noted that intensive and diverse investigated terraneous mollusks of Uzbekistan have been held mainly for the past 15-20 years, as a result, a large amount of material has been accumulated that has not yet been generalized. Having studied in detail the vertical distribution of terraneous molluscs, since knowledge of the distribution of mollusks in various types of biotopes and their vertical placement can serve as the main basis for the development of measures to

* Corresponding author: feruz.bio@yandex.ru

combat mollusks that are pests of crops and intermediate hosts of helminths, and preventive measures against them.

As we know, a number of terraneous mollusk species are intermediate and reservoir hosts of cattle helminth and pests of agricultural plants, however, the number of species of economic importance is unknown to this day. Therefore, the fourth tasks of the article are to determine the number of species - intermediate hosts of helminths and to revise the composition of intermediate hosts of a number of nematodes of the researched territory and to find out the role of new and little-known species of terraneous mollusks in the life cycles of helminths.

2 Materials and methods

The actual basis of the article is based on the materials of our collecting carried out in the mountain ranges: Karzhantau, Ugham, Pskem, Chatkal, Talas, Kuramin, Ferghana, Alai, Turkestan, Zarafshan, Gissar, Nuratinsky, Kugitangtau, Baysuntau, Babatag and in Ferghana, Babatag and in the Ferghana, Chirchik-Akhangaran, Zarafshan, Kashkadarya and Surkhan-Sherabad valleys, as well as extensive collections of the Zoological Institute of the Academy of Sciences of Russia, the Zoological Museum of Moscow University.

3 Results and Discussion

In addition, the collections of Samarkand University, the Institute of Zoology and Parasitology of the Academy of Sciences of the Republic of Tajikistan were partially used. The researches have shown that the faunal spectrum of terraneous mollusks of Uzbekistan and bordering territories is very diverse (Table 1). The family Cochlicopidae in the research area is represented by one genus (*Cochlicopa*) and 8 species, which is 4.67% of malacofauna. Representatives of this family are widespread in the Palearctic. The following locations were found in the investigated territories. All species live in interzonal biotopes.

The family Orculidae in the research area is represented by 2 subfamilies: *Orculinae* *Lauriinae*, the range of which covers the Mediterranean countries, the Canary and Azores Islands, Europe, Crimea, the Caucasus, Central Asia. Subfamily *Orculinae* is represented by one monotypic genus - *Sphyradium* (range, as in the family as a whole). *S. doliolum* was found in our area, living in a large set of biotopes.

Lauriinae also consists of one genus and the species *Lauriacylindracea*. According to A.A. Shileiko (1984), its range covers the Primorsk regions of Europe from the islands of Bergholm and southern Norway along the Atlantic and Mediterranean coasts to Asia Minor, North Africa, Crimea, Transcaucasia, Dagestan and Kopetdag.

They were first discovered by us in Central Asia on the Kugitangtau peninsula. The *Valloniidae* family is very extensive, known in the Palearctic.

There are 2 subfamilies in the research area: *Acanthinulinae* (with 1 genus and species) and *Valloniinae* (with 1 genus and 3 species). Representatives of the genus *Vallonia* live in sufficiently moistened biotopes.

The subfamily *Acanthinulinae*. *Acanthinulaaculeata* in the researched area was first discovered on the Kugitangtau ridge. The natural range of this species includes Europe, North Africa, Crimea, the North Caucasus, Transcaucasia.

The *Pupillidae* family in the fauna of the territory under consideration is represented by 11 species (which is 6.43% of the malacofauna) belonging to 2 genera. The genus *Gibbulinopsis* with an area including the Caucasus, Central and Central Asia, the Far East and Japan.

3 species were found in the researched area, 2 of them (*Gibbulinopsisgracilis*, *G. nanosignata*) are endemic, known in the north-western part of the Pamir-Alai (Hissar, Turkestan ridges). Representatives of the genera *Gibbulinopsis* are very drought-resistant, they live in steppe, semi-desert and mountain steppe areas, where they are kept in small-block scree and dead plant remains.

Table 1. Faunal spectrum of terraneous mollusks of Uzbekistan and bordering territories.

Family of molluscs	Quantity		%	Number of types	%
	subfamilies	genus			
Cochlicopidae	-	1	2.22	8	4.65
Orculoidae	2	2	4.44	2	1.16
Valloniidae	2	2	4.44	4	2.32
Pupillidae	1	2	4.44	11	6.39
Vertiginidae	2	3	6.66	7	4.06
Chondrinidae	-	1	2.22	1	0.58
Puramidulidae	-	1	2.22	1	0.58
Buliminidae	1	8	17.77	53	30.81
Vitrindae	-	1	2.22	1	0.58
Bradybaenidae	-	2	4.44	17	9.88
Hygromiidae	5	12	26.66	38	22.09
Agriolimacidae	-	2	4.44	7	4.06
Limacidae	-	1	2.22	4	2.32
Parmacellidae	-	1	2.22	7	4.06
Ariophontidae	-	1	2.22	4	2.32
Gastrodontidae	-	1	2.22	1	0.58
Succinidae	-	4	8.88	6	3.48

The genus *Pupillae* of the studied territory is represented the fauna by 8 species, of which *P. muscorum* is Holarctic. The range of *P. triplicata*, *P. bigranata*, *P. starii* covers southern, central-western Europe, the North Caucasus and Transcaucasia, Asia Minor and Near East and mountainous areas of Central Asia. The ranges of *P. gallae*, *P. turcmenica* are limited to several mountain ranges. *P. anzobica* is found only on the Hissar ridge.

The range of the *Truncatellinidae* subfamily covers all continents except Australia, represented by 9 genera, in Central Asia - 2 genera (*Columella*, *Truncatellina*) and 5 species.

The genus *Columella* is Holarctic, includes 5 species. 3 species were found in the research area: *C. columella*, *C. edentula*, characterized by an extensive range, and *C. intermedia* - endemic to the Tien Shan. It was first discovered by us on the Pskov and Chatkal ridges. The range of the genus *Truncatellina* includes Europe, the Caucasus, Eastern, Southern Africa, Central Asia, and is represented in the rese region by 2 species living in dry plant remains on open slopes and in woodlands.

The family *Chondrinidae* in the studied territory is represented by one genus *Chondrina*, whose natural range includes Southern, Central Europe, North-West Africa, Crimea, the Caucasus, Near Asia, Iran, Kopetdag. It was first discovered by us on the Kugitangtau ridge.

The family *Buliminidae* in the malacofauna of Uzbekistan and adjacent territories is distinguished by the diversity and richness of species. It is represented here by 8 genus and 52 species of mollusks (30.40% of the malacofauna).

The area of the *Pseudonapaeinae* subfamily. Its range includes Transcaucasia, Iran, Afghanistan, India, Central and Central Asia. The total number of genus is at least 20, 6 genera, 45 species live in the research area, which is 26.31% of the malacofauna.

The genus *Ottarosenia* with the only species *O. varenzovi* is endemic to Kopetdag, first discovered on the Kugitangtau ridge.

The genus *Pseudonapaeus* in the malacofauna of Uzbekistan and neighboring territories is distinguished by the variety and richness of species. It is represented by 31 species of mollusks, which is 18.12% of malacofauna. Among them there are endemics with a narrow range, confined to certain mountain ranges. This is *Ps. shahristanicus*, characteristic of Turkestan, *Ps. zaravshanicus* - for Zarafshan, *Ps. kasnakovi*, *Ps. otostomus* - for Gissars, *Ps. chodschendicus* - for Kuraminsk, *Ps. naukaticus* - for Alai, *Ps. goldfussi*, *Ps. arislanbobica* - to the Ferghana ranges. The wider range is distinguished by *Ps. sogdiana*, *Ps. eremita*, common in Iran, Afghanistan, India, Kopetdag, on the Alai and Turkestan ranges. The range of the *Turanena* genus includes Transcaucasia, Northern Iran, Kopetdag, Himalayas, Pamir-Alai, Tien Shan, Western China. Ten species, accounting for 5.84% of the malacofauna, are found in the studied area. *T. scalaris* and *T. herzi*, which are widespread in Northwestern Iran, Armenia, and Mount Elbrus, have a wider range. In the studied area (Babatag, Baysuntau) were found for the first time. Narrow-areal endemics include *T. cognata*, *T. stschukini*, distributed only in the Western Tien Shan. There are species confined to certain mountain ranges. This is *T. meshkovi*, characteristic of the Pskem Range.

The genus *Subzebrinus*, with a single species *S. labiellus*, occurs only in the Tarbagatai, Zailisk and Dzungarian ranges. We first discovered it in the Western Tien Shan (in the Ugam and Pskem Ranges). *Mastoides* is an autochthonous genus. Two species (*M. orloffensis*, *M. albocostatus*), which are narrow-areal endemics of the Fergana Range, are represented in the studied area.

The genus *Laevozebrinus* in the fauna of the studied area is represented by 2 species. Both species are endemics: *L. ujfalvyanus* (Chatkal and Fergana Ranges) and *L. lenis* (Karzhantau Ridge).

The subfamily Chondrulopsinae is represented by two genera - *Chondrulopsina* (its areal covers mountainous areas of Central Asia, northern Afghanistan, western China) and *Siraphoroides* (known only on the Fergana Ridge).

The family Vitrinidae in the studied territory is represented by a single genus and species *Phenacolimnax annularis*, whose range covers Europe, North Asia, Central Asia, is widely distributed in the mountain ranges of Karzhantau, Ugam, Pskov, Hissar, Baysuntau.

Family Bradybaenidae. The greatest diversity is observed in South-east Asia, in addition, the range of this family covers the Philippine Islands, Japan, some species penetrate into Europe. Two genera are represented in the Bradybaenidae in the studied territory: *Bradybaena* (younger), *Ponsadenia* (older). Genus *Ponsadenia*. Its range covers the Tien Shan, the Dzungarian, Tarbagatai and Western China ranges. On the investigated territory there are 2 species: *P. duplocincta* and *P. semenovi*. Of these, *P. semenovi* is distributed outside the studied territory - in Western China. Both species differ in their ecological characteristics. *P. durliosinista* is a hygrophilous species, found in mountainous areas, inhabiting forested areas. *P. semenovi* can be found in all landscape-geographical zones. The range of the genus *Bradybaena* covers Central Asia, all of East Asia, north to Kamchatka, the Philippines and Japan. Two species inhabit Europe. There are 14 species in the studied area, which can be called endemics of the Tien Shan. *Br. lantzi*, *Br. almaatini*, *Br. cavimargocavimargo* have a wider areal: covers the mountain systems of Tien Shan, Pamir-Alai. The remaining species are confined to certain mountain ranges. These are *Br. dichrozona*, *Br. saturata* - Chatkal and Kuramin ranges. All species of this genus differ in their ecological features: *Br. phaeozona*, *Br. almaatini*, *Br. dichrozona*, *Br. saturata* inhabit among shrubs, on rubbly areas, *Br. reglucens*, *Br. stoliczkana*, *Br. fedtschenkoi* - under stones, are also confined to rocky talus. The Hygromiidae family in the malacofauna of Uzbekistan and adjacent territories is distinguished by the diversity and richness of species

(after this. *Buliminidae*). Here it is represented by 5 subfamilies, 12 genera and 38 species, which make up 22.22% of the malacofauna.

The *Hygromiidae* family in the malacofauna of Uzbekistan and bordering territories is distinguished by the diversity and richness of species (after this. *Buliminidae*). Here it is represented by 5 subfamilies, 12 genera and 38 species, which make up 22.22% of the malacofauna.

The *Trichiinae* subfamily covers Europe, the Mediterranean, the Caucasus and Central Asia, is represented by 5 genera, of which 3 genera (18 species) are endemic in Central Asia.

The genus *Nanaja* - endemic to Western Tien Shan - is represented by 2 species: *N. cumulata*, *N. illuminata*. Both species are narrowly localized until discovered in the Pskem Range. They live mainly in large-fragmented sedentary showers, not far from streams. Both species are narrowly localized, so far found in the Pskem Range. They inhabit mostly in coarse-clastic low-mobile screes, not far from streams.

The genus *Odontotermia* is an endemic of the Chatkal and Fergana Ranges and is represented by two species: *O. diplodon* and *O. monodon*. Both species occur in mountain areas at an altitude of 2,500-2,800 m above sea level. They live among shrubs and deciduous forests; they prefer wetter deciduous forests.

The genus *Leucozonella* is endemic to Central Asia; 14 species are currently known in the study area. The most widespread are *Leucozonellacarauodes*, *L. rubens*, *L. mesoleuca*, and *L. rufispira*, which inhabit Pamir-Alai and Tien Shan. The remaining species are endemics with a limited range: *L. ferghanica*, *L. crassicosta*, *L. hypophaea*, *L. angulata* - Western Tien Shan; *L. schileykoi* - only the basin of the Ak-su River in the Turkestan Range, and *L. globulifopmus* - in the Alai Range. All species of this genus differ in their ecological features: *L. ferghanica*, *L. angulata* prefer wetter biotopes, not far from spring waters; *L. mesoleuca*, *L. caria* inhabit among bushes, in open areas; *L. rufispira*, *L. reterri* are confined to coarse clastic rocky screes; *L. hypophaea*, *L. schileykoi* - to dry open rocky slopes.

The range of the genus *Helicopsis* includes Middle and Southern Europe, Crimea, Transcaucasia, Asia Minor, North Africa, Central Asia (Kopetdag, Kugitangtau). The only species in the studied area is *H. likharivi*, which inhabits open rocky slopes and was first discovered on the Kugitangtau Ridge.

The range of the genus *Xceropicta* covers the Caucasus, Crimea, in some places along the northern shore of the Black Sea, the Balkan Peninsula, North-East Africa, Central Asia. Two species are known in the studied area: *X. candaharica* - with a wide area and *X. kupiskii* - distributed in places in the Chatkal, Alai, and Nurata Ranges. Both species are xerophilous.

Endemic subfamily. *Archaicinae* includes 4 genera. Its area of distribution is limited to mountainous areas of Central Asia. The range of the genus *Archaica* includes the Western, Southern Tien Shan and Alai ridge.

6 species were found in the investigated territory. The wide range is distinguished by *A. heptapotamica* - from the northwestern part of the Ferghana and the central part of the Talas ranges to the west to the westernmost spurs of the Tien Shan, the western part of the Kyrgyz range. The remaining species are endemic with a narrow range. *A. apollinsis* was found on the Ferghana and Alai, *A. papanica* - Alai, *A. karjantauca* - Karzhantau, *A. eleorina* - on the Zarafshan ridges. In ecological terms, representatives of this genus do not differ. In ecological terms, representatives of this genus do not differ. They live mainly in large - fragmented showers with high herbage.

The genus *Hissarica* (monotypic) with a single species *H. incltatus* is known only on the Zarafshan ridge. *Leucarchaica* is also a monotypic genus, represented by a single species *L. rudimentifera*, known only on the Fergana Ridge. The *Euomphaliinae* subfamily in the

research area includes the only (imported) genus *Monacha* and the species *M. carthusiana*, found on the northern slopes of the Turkestan ridge.

The range of the Paedhoplitinae subfamily is limited to the mountainous regions of Central Asia, its representatives originated locally from some Trichiinae, regardless of the European-Caucasian Hygromiinae and Euomphaliinae. It is represented by 2 genera - *Angomphalia* and *Paedhoplita*.

The genus *Angiomphalia* includes 5 species. Of these, *A. regeliana* is widely distributed across the Tien Shan and the northern spurs of the Pamirs. It lives in various biotopes, in flat zones - among thickets of grasses, near spring waters; in mountainous areas - among shrubs, prefers the northern slopes of hills.

The ranges of other species are narrow and narrowly localized. *A. copisa* is found only on the Chatkal ridge, *A. exasperata*, *A. calestomontana*, *A. seductilis* - on the Central and Northern Tien Shan. It was first discovered by us in the northeastern part of the Ferghana Ridge.

The genus *Paedhoplita* in the research area is represented by 2 species: *P. lentina* - on the Ferghana and Chatkal ridges; *P. buamica* - in the eastern part of the Kyrgyz and northwestern part of the Ferghana ridges.

The range of the Agriolimacidae family. Its range covers almost the entire Holarctic, but most genera and species are characteristic of Mediterranean and coastal countries. In general, the family is represented by 6 genera. 2 - *Deroceras*, *Lutopelte* were found in the research area.

The range of the genus *Deroceras* includes most of the northern hemisphere.

Most species live in the Mediterranean subdistrict. There are 6 species in the research area. Inhabitants of very moist biotopes are often found along the shores of reservoirs.

The genus *Lutopelte* with a single species *L. maculata* in the research area was found on the southwestern slopes of the Fergana Ridge, in the valley of the Zarafshan River. It lives among plants, under rocks, as well as among shrubs, along mountain rivers.

The range of the Limacidae family. Its range covers Europe, the Mediterranean region of North Africa, the Caucasus, Near and Central Asia. It is represented by a total of 10 genera. Representatives of the endemic genus *Turcolimax* with 4 species are common in the research area. The wider range is distinguished by *T. natalianus*, common in Northern Tien Shan, Dzungaria, Tarbagatai and Western China. The range of *T. turkestanus* covers the Northern and Central Tien Shan. *T. ferghanus* is characterized by a narrow range – only the Chatkal ridge.

The range of the Parmacellidae family, according to I.M. Likharev and A.I. Victor (1980), consists of two parts isolated from each other: a) North Africa (from Alexandria to Morocco) together with Portugal and southern Spain; b) eastern Transcaucasia, Northern Iran, Kopetdag, Central Asia, Afghanistan. It is represented by 2 genera: *Parmacella* and *Candacharia*. Representatives of the genus *Candacharia* are distributed in the research area, whose range covers the mountain systems of the Tien Shan, Pamir-Alai, mountainous regions of Afghanistan and North-eastern Iran.

The genus *Candacharia* is represented by 7 species, 2 of which - *C. izzatullai*, *C. langarika* - are endemic with narrow ranges confined to certain mountain ranges (Zarafshan).

The family Ariophantidae. Most of its species, according to K. K. Uvalieva (1990), live in the tropics.

In Central Asia, it is represented by a single genus - *Macrochlamys*. There are 4 species in the research area. Among them, the most widely developed. *schmidtii* – in all mountain systems of Central Asia. *M. turanica*, *M. sogdiana* are common in the Tien Shan and Pamir-Alai mountain systems. *M. kasnakovi* does not go beyond the Pamir-Alai.

The Gastrodontidae family. The main part of its species lives in the Western Palearctic. In the research area, the genus *Zonitoides* is represented by the only species, *Z. nitidus*, found in very moist biotopes.

The family Succineidae. The main number of its species also lives in the western part of the Palearctic, 4 genera are represented in the fauna of Central Asia.

The range of the genus *Novisuccinea* covers Central and Eastern Asia to the north – Kamchatka, the Philippines. 2 species were found in the research area: *Novisuccineaevoluta*, *N.martensiana*, widely distributed in Tien Shan, Altai, Transbaikalia, Tibet. The most characteristic habitats for these species are wet rocks near mountain rivers and streams, waterfalls, as well as alpine meadows, where they adhere to scree and rock outcrops.

The genus *Pamirsuccinea* is represented by a single species, *P. eximia*, distributed in Pamir-Alai. It lives near water in humid biotopes. The area of genus *Oxyloma* covers Europe, Caucasus and Central Asia. Two species of this genus inhabit the area of researches. *O. elegans* in the study area is found locally in the Turkestan, Alay, Fergana, Chatkal, and Alay Ranges in damp places near water, often among coastal vegetation. *O. sarsi* is widespread in Transcaucasia, Western Siberia, and Central Asia and inhabits the most humid biotopes.

The range of the genus *Succinea* includes Europe, the Caucasus, Altai, Siberia, and Central Asia. In general, the genus is represented by 3 species. *Succinea putris* inhabits in area of researches, in humid habitats, but excessive moisture is avoided, inhabits also flooded high grass meadows.

4 Conclusion

Having studied the faunistic spectrum of terrestrial mollusks in Uzbekistan and adjacent territories, the ecological groups of terrestrial mollusks in relation to the moisture factor and habitats have been determined. As a result, the data on the species composition of terrestrial molluscs have been significantly replenished by including a number of species new to the fauna of Central Asia; the boundaries of the ranges of a number of species have been clarified; revised taxonomic position of some forms; the originality of the malacofaunas of different altitudinal belts was revealed.

References

1. Z. Izzatullaev, Little-known terrestrial mollusks (Mollusca Gastropoda) of the fauna of Central Asia, *Izv. AN Taj SSR. Dept. biol. sciences*, **2**, 39-44 (1975)
2. V.A. Lindholm, Description of two types p, *Buliminus* from Russian Central Asia, *Journal of Annual Zool. Mus.*, **23**, 304-320 (1922)
3. V.A. Lindholm, Mollusks: Pamir, expeditions, **8**, 29-64 (1928)
4. I.M. Likharev, E.S. Rammelmeyer, Terrestrial mollusks of the fauna of the USSR. Key to the fauna of the USSR (Nauka, Moscow-Leningrad, 1952)
5. I.M. Likharev, On the systematic position of some Central Asian terrestrial mollusks, *Tr. Zool. Institute of the USSR Academy of Sciences*, **21**, 179-185 (1955)
6. P.V. Matekin, Materials on the fauna of terrestrial mollusks of Central Asia: Dis doct. biol. sciences, 516 (1960)
7. P.V. Matekin, Keys to the shell mollusks of Central Asia - intermediate hosts of helminths: Helminths of animals in Kyrgyzstan and adjacent territories, Frunze, 97-136 (1966)

8. Ya.I. Starobogatov, Eurasian species of the genus *Cochlicopa* (Gastropoda, Pulmonata, Cochlicopidae), *Ruthenica*, **5**, **2**, 105-131 (1996)
9. K.K. Uvalieva, Terrestrial molluscs of Southern Altai and neighboring mountainous regions: Author's abstract. dis. ...cand. Biological Sciences, 19 (1965)
10. A.A. Shileiko, The structure and systematic affiliation of species of the genus *Siraphorus* Lindholm, 1925 (Gastropoda, Enidae), *Scientific Dokl. Higher School of Biological Science*, **9**, 40-46 (1977)
11. A.A. Shileiko, Terrestrial mollusks of the superfamily Hellicoidae, *Fauna of the USSR. shellfish*, Science Leningrad branch, **3**, **6**, 384 (1978)
12. A.A. Shileiko, Research of type species of some taxa of the genus group in the family Buliminidae (Enidae, Gastropoda) species of Central and Central Asia, *Zool.zhurn*, **57**, **3**, 344-358 (1978)
13. A. Pazilov, F. Gaibnazarova, M. Saidov, Regularities of vertical distribution of terrestrial mollusks in Uzbekistan and adjacent territories (Fan, Tashkent, 2014)
14. F. Gaibnazarova, Character of variability of signs of the reproductive apparatus *Pseudonapaeus Albiplicata* From Chatkal, Kuraminsky Ridges-Guliston Davlat Universiteti Axborotnomasi, 3 (2015)
15. F. Gaibnazarova, A. Pazilov, *Conchological variability of terrestrial mollusks Gibbulinopsis nanosignata of the Turkestan and Zarafshan ranges*, Zoological studies of the regions of Russia and adjacent territories, materials of the III international scientific conference, Nizhny Novgorod (2014)
16. A. Pazilov, F. Gaibnazarova, Kh. Karimova, Terrestrial mollusk complexes in various biotopes in zarafshan range, *Journal NX- A Multidisciplinary Peer Reviewed Journal* ISSN: 2581-4230 (2020)
17. F. Gaibnazarova, Kh. Karimova, Z. Muhammadiyev, "Geographical and ecological analysis of dry mollusks in Uzbekistan and adjacent regions" (2021)
18. F. Gaibnazarova, Character of variability of signs of the reproductive apparatus of *Pseudonapaeus albiplicata* from the Chatkal and Kuramin ridges, *Biological sciences of Kazakhstan*, 3 (2014)
19. M.A. Marinin, M.A. Karasev, G.B. Posphehov, A.A. Pomortseva, V.N. Kondakova, V.I. Sushkova, Comprehensive study of filtration properties of pelletized sandy clay ores and filtration modes in the heap leaching stack, *Journal of Mining Institute*, **259**, 30-40 (2023)