

Optimizing Food Security and Environmental Sustainability via Agroecology and Sustainable Intensification Strategies

Vijilius Helena Raj^{1*}, S Vinod Kumar², M. Sabir³, Dinesh Kumar Yadav⁴, Shekhar Sharma⁵, Haider Alabdeli⁶, Vanya Arun⁷

¹Department of Applied Sciences, New Horizon College of Engineering, Bangalore

²Institute of Aeronautical Engineering, Dundigal, Hyderabad

³Department of English, Nagpur Institute of Technology, Nagpur, India

⁴Lloyd Institute of Engineering & Technology, Knowledge Park II, Greater Noida, Uttar Pradesh 201306

⁵Lloyd Institute of Management and Technology, Greater Noida, Uttar Pradesh, India-201306

⁶The Islamic University, Najaf, Iraq.

⁷Department of Electronics and Communication Engineering, IILM University, Greater Noida

*Corresponding author: vijilius_23@gmail.com

Abstract. The science of ecology is incorporated into farm development and operation through agroecological techniques. A paradigm shift in agriculture is essential to combat hunger, adapt to climate trade, and mitigate environmental degradation. By doing this, researchers may further acknowledge the interdependence of farmed and nonfarmed landscapes and the variety of products and services that robust ecosystems offer, including resilience, nutrient cycling, and pest control, all of which can help sustain yields. Agro-ecology relies heavily on the knowledge and experience of farmers since it fosters independence and decreases reliance on costly outside resources. The concepts of sustainable intensification and agroecology are examined in this paper as additional strategies to address the global issue of increasing food production while lowering environmental impacts. Also, this study evaluates how effectively these approaches boost crop yields, lower environmental costs, and build resilience to climate unpredictability by closely examining existing programs, integrated management strategies, and field experiments. Supporting the findings is a comparative table that shows several techniques of sustainable intensification and how they have an effect on yields and environmental costs. In order to expand resilient, sustainable, and equitable food systems, a discussion of the necessity of a paradigm shift towards agroecological strategies is addressed in the paper's conclusion.

Keywords:- Food production, Agroecology, Crop yield, Sustainable intensification, Environmental impacts, Integrated management practices.

1 Introduction

Sustainable intensification that is getting increasingly reputation, is a key tactic for tackling the troubles introduced up via the growing international population, climate trade, worries approximately food safety, and the requirement of resource conservation in agriculture [1]. Environmental stewardship, which entails the preservation and regeneration of herbal capital and ecosystem offerings, should be balanced with this, even though it regularly method higher output via extended productivity and generation use. This shows addressing intake patterns, reducing waste, protecting biodiversity, and optimizing aid usage to be able to ensure that sustainable manufacturing levels meet human requirements [2]. On the same time, "agroecology" is gaining recognition as a means of incorporating ecological ideas into the planning and management of agricultural systems to strike a balance among environmental, productivity, and sustainability objectives. Agroecology places an emphasis on ecological and statistics intensification further to technological improvements inside the context of sustainable intensification [3]. A paradigm shift in agriculture is essential to combat hunger, adapt to climate trade, and mitigate environmental degradation. This has been recognized by stakeholders from governments, academia, international organizations, non-governmental organizations, and the personal area increasingly, and has sparked discussions amongst them. Fig. 1 shows how the sustainable intensification deals when the resources including labour, wages, etc. Furthermore, livestock farming structures are scrutinized due to concerns approximately animal welfare, the effects they have on the environment, and the health risks linked with consuming an excessive amount of meat, even if they greatly contribute to protein-rich diets, jobs, and the preservation of cultural heritage [4].




Fig. 1: Sustainable intensification while dealing with limited resources [3]

Agroecology uses insights from natural ecosystems to create a strong basis for sustainable intensification. Organic and ecological methods such as allelopathy, competition, nitrogen fixation, nutrient cycling, and soil revitalization are investigated in detail and protected into this method. Agro ecology also aims to reduce reliance on non-renewable inputs that jeopardize consumer health, the environment, and farmers' livelihoods. Agro ecology relies heavily on the knowledge and experience of farmers since it fosters independence and decreases reliance on costly outside resources. Agro ecology also welcomes group intelligence to efficiently solve shared resource and agricultural concerns. Fundamental ideas from natural ecosystems are crucial, such as preserving perennial vegetative cover or using appropriate substitutes like cover crops and intercropping. By maintaining an organic matter supply that is steady, these methods preserve soil health, save water, and support soil biological activity. Furthermore, agroecology prioritizes crop diversity and animal husbandry to naturally manage pests and diseases and foster diversity at both field and landscape scales. To replicate the diversity of steppe or prairie vegetation, for example, crop rotations can include combinations of perennial grasses. To successfully integrate these ideas into existing farming methods, however, will necessitate structural changes in the agricultural industry.

Intensification is often given precedence over agroecological methods when it comes to increasing agricultural productivity. Agroecology provides a broad paradigm shift, in contrast to the narrow focus of organic agriculture and sustainable intensification programs. It questions the idea that small changes to the existing agroindustrial system, sometimes referred to as "greening," will adequately address agricultural sustainability. Reducing input consumption is a good start in the right direction in terms of technology, but it is not enough to make the shift to a more independent and self-sufficient farming system. The sustainability of monoculture and the excessive reliance on outside inputs have been called into question by a true shift toward agroecology. Rather, it looks to traditional agricultural systems as role models that run without heavy use of agrochemicals, prioritize biodiversity, and sustain annual yields that are constant.

2 Principles of Agroecology

Agroecology has many different meanings and interpretations. In 2019, 197 countries unanimously adopted the 10 elements of agroecology to provide guidance for the Food and Agriculture Organization (FAO) in realizing its vision of agroecology. those elements are the outcome of a collaborative approach in which the FAO invited other parties to participate. furthermore, in that same year, the high-level Panel of experts on food security and nutrition (HLPE) published the 13 agroecological principles. Serving as a crucial link among scientific knowledge and policymaking, the HLPE, beneath the path of the UN Committee on international food security (CFS), conducts in-depth, evidence-based assessments and fosters policy discussions. Agroecological techniques and other improvements for sustainable agriculture and food systems that increase meals protection and nutrition was a paper published with the aid of the HLPE in 2017 in reaction to a request from the CFS. This illustrates how agroecology is starting to be acknowledged as a viable remedy for international issues.

For the purpose of developing a sustainable food system, the HLPE provided three guiding principles. Though there are notable synergies between these principles that facilitate the identification of mutually beneficial solutions, it is acknowledged that trade-offs between them may not always be avoidable in specific situations. To increase productive efficiency and encourage investment, for example, one can increase resilience to animal diseases, climate change, and price variations (HLPE, 2016). The necessity of implementing a comprehensive strategy for food systems is shown by the connections between these operational concepts. These operational strategies are presented in Fig. 2.




Fig. 2: Operational principles of agro-economy

- a) **Improve Resource Efficiency**- Improving Resource Efficiency aims to optimize resource use for economic gains, environmental sustainability, and social equity among smallholders, pastoralists, and family farmers. Production can be increased by the use of both new and existing agricultural technology and practices to fill yield gaps. This can have a favorable impact on the production of animal feed and animals. For vulnerable groups, this approach may improve economic situations, diets, and nutrition [9]. Agroecological resource management systems are made up of interdependent parts that work together to collect, structure, and disseminate operational data related to precision farming. While rich countries have long embraced precision farming, developing nations like Brazil, China, India, and several African countries are now seeing an increase in the practice. A number of prototype and commercial sensor platforms intended for agricultural applications were recently detailed in international research journals [9, 10].
- b) **Strengthen Resilience**- Resilience building involves encouraging effective resource use while strengthening the ability to endure shocks and bounce back from them. In order to guarantee sustainable resource consumption even under challenging circumstances, this includes efforts targeted at strengthening adaptive capacities and controlling risks [11]. Modern agricultural systems can benefit from the many ideas and methods from traditional farming systems in order to strengthen their resistance to climate-related problems. Reducing vulnerability to climate variability is a key function of several of these agroecological techniques, including crop diversification, harvesting, organic soil management, maintaining local genetic diversity, and integrating animals. The resilience of traditional agroecosystems can be developed using the agroecological traits that underpin them as a model for developing resilient agricultural systems, therefore understanding these traits is crucial. The evaluation of agricultural performance following extreme weather events, such as droughts and hurricanes, over the past 20 years has brought attention to the significant relationship between farms with higher biodiversity and their ability to withstand climate disasters [12].
- c) **Secure Social Equity / Responsibility**- promoting fairness and accountability while addressing social and ethical challenges within societies is a key aspect of ensuring secure social equity and responsibility. This consists of things like gender equality, social safety, human rights, income fairness, and cultural preservation. It calls on people, organizations, and businesses to fulfill their obligations to protect the environment, enhance human welfare, and enhance animal welfare. Acknowledging the diversity of norms and behaviors among nations and communities, achieving this goal requires a variety of context-specific techniques [13].

Offering women and other historically marginalized stakeholders the chance to lead the movement is essential if agroecology is to fulfill its dual goals of promoting ecological and social well-being. Agroecology's social and gender implications go beyond its explicitly social features and are fundamental to all facets of the field [14]. This emphasizes how crucial it is to incorporate social justice and gender equality into all facets of agroecological practices and policies.

3 Sustainable Intensification

Providing food for the world's growing population while also minimizing environmental damage is a major concern facing civilization today. Recent initiatives support "sustainable intensification" as a workable land use strategy, backed by institutions like the United Nations and the US Agency for International Development. Protecting ecological services and biodiversity while increasing food production is the goal. For those tackling this crucial issue, there remains uncertainty regarding the exact meaning and consequences of sustainable intensification [15, 16]. In the field of agriculture, it is extremely difficult to increase yields while also reducing environmental effects in order to ensure global food security, Fig. 3 shows the role of agr-ecosystem for sustainable intensification to reduce the concern of maximizing yields. The adoption of integrated soil-crop system management strategies, grounded in modern crop ecophysiology and soil

biogeochemistry, has shown promising results. These techniques considerably increased the average yields of rice, wheat, and maize without necessitating the use of nitrogen fertilizer, according to a study [17]. Particularly, the yields of 10.5 mg/ha to 8.9 mg/ha, 8.5 mg/ha to 14.2 mg/ha, and wheat, rice, and maize increased from 7.2 mg/ha to 8.9 mg/ha, respectively. Life-cycle analyses and model simulations that demonstrate notable reductions in greenhouse gas emissions and reactive nitrogen losses further underscore the benefits of this approach. By scaling up these strategies to achieve 80% of observed yields by 2030 across China's 2012 planting area, the need for both direct human consumption and expanding animal feed requirements might be efficiently supplied while simultaneously limiting the environmental impact of intensive agriculture [17]. Overuse of nitrogen fertilization in China's largely agriculturalized areas has resulted in major environmental issues, including air, soil, and water enrichment with reactive nitrogen from agricultural sources. The study outlined in [18] examines grain yields and nitrogen loss patterns in two intensive double-cropping systems in China: waterlogged rice/upland wheat in the Taihu region and rainfed maize and irrigated wheat on the North China Plain. Current nitrogen application approaches, averaging 550–600 kg per hectare every year, do not result in appreciable gains in crop yields, in contrast to optimal nitrogen fertilization methods that have the potential to save 30–60% of nitrogen usage. As an alternative, they cause the environment to lose nitrogen by around twice as much. In wheat/maize systems, the main sources of nitrogen loss are nitrate leaching and ammonia volatilization, which account for 19% and 24% of nitrogen loss, respectively. These losses are made worse by elements including calcareous soils and intense summer rainfall. Cost-effective methods for lowering greenhouse gas (GHG) emissions are becoming more and more in demand as attempts to combat climate change increase. Given that it is a significant emitter, agriculture presents options for mitigation as well [19].




Fig. 3: Consistent increasing of agroecosystem services

The study conducted in [19] evaluates the overall impact of historical agricultural intensification (1961-2005) on greenhouse gas (GHG) emissions. Higher yields have reduced emissions by up to 161 gigatons of carbon (GtC), or 590 gigatons of carbon dioxide equivalent (GtCO₂e), since 1961, even in spite of the increasing emissions linked to activities like fertilizer manufacturing. Compared to 1961 technology, there has been a reduction of 68 kilograms of carbon (kgC), or 249 kilograms of carbon dioxide equivalent (kgCO₂e), emissions for every dollar invested in agricultural yields. This means that 3.6 GtC, or 13.1 GtCO₂e, of emissions have been avoided annually. An evaluation was conducted in a field experiment to determine how well Integrated Nutrient Management (INM) reduced nitrogen (N) losses and enhanced wheat (*Triticum aestivum* L.) growth and production metrics [20]. The application of NPK + Azotobacter + mycorrhiza (Tram) promoted microbial activity, root growth, urease activity, and nitrogen utilization efficiency, and improved soil nitrogen dynamics when compared to plots receiving only NPK. It also increased the availability of phosphorus (P) and potassium (K). Plots treated with INM demonstrated reduced denitrification losses (2.4-2.5 kg/ha) and cumulative ammonia flow (10.2-10.6 kg/ha) as compared to those treated with NPK alone. The INM plot that received Tram had the maximum yield (6.4 t/ha). Meanwhile, studies described in [21] evaluated the effects on greenhouse gas emissions from China's production of staple crops of integrating biomass pyrolysis and electricity generation with methane and nitrogen mitigation strategies using a thorough life-cycle assessment technique. The findings showed that annual emissions decreased significantly from 666.5 to -37.9 Tg CO₂-equal. The usage of biochar, which trapped carbon, and the switch to bioenergy from fossil fuels become the number one reason of this decline. The incorporated system's 8.3% increase in crop output, 25.5% lower in reactive nitrogen losses, and 125–2,483 Gg yearly reduction in air pollution emissions also contributed to a 36.2% advantage in net environmental and economic advantages. The comparison of previous conducted studies are mentioned in Table 1, with respect to their purpose, techniques adopted and impact on the technique in enhancing the yields.

Table 1: Comparative analysis of different methods contributing in sustainable intensification

Reference	Purpose	Methodology	Impact on Yields	Impact on Environmental Costs
-----------	---------	-------------	------------------	-------------------------------

[17]	Increase in yields for wheat, rice and maize without additional nitrogen fertilizer	Crop ecophysiology and soil biogeochemistry based Integrated soil-crop system management	Rice: 7.2 to 8.5 Mg/ha, Wheat: 7.2 to 8.9 Mg/ha, Maize: 10.5 to 14.2 Mg/ha	Reduced reactive nitrogen losses and greenhouse gas emissions
[18]	Analysis of nitrogen loss pathways in double-cropping systems	Comparison of grain yields and nitrogen losses in waterlogged rice/upland wheat vs. irrigated wheat/rainfed maize systems	No significant yield increase with high nitrogen use	Optimal nitrogen fertilization can save 30-60% nitrogen with reduced losses
[19]	Evaluation of GHG emissions from agricultural intensification	Historical data analysis (1961-2005)	Adoption of higher yields prevented emissions	Prevented emissions of up to 161 GtC, equivalent to 590 GtCO ₂ e
[20]	Impact of Integrated Nutrient Management (INM) on wheat yield and N losses	Field experiment using NPK + Azotobacter + mycorrhiza vs. NPK alone	Highest yield with INM: 6.4 t/ha	Lower ammonia and denitrification losses with INM
[21]	Reduction of GHG emissions through integration of biomass pyrolysis and electricity generation	Life-cycle assessment of integrated system for staple crop production	8.3% increase in crop yield	Significant emissions reduction from 666.5 to -37.9 Tg CO ₂ -equivalent per year, plus net environmental and economic benefits increase

4 Benefits of Agroecology and Sustainable Intensification

Agroecology and sustainable intensification offer numerous benefits as presented in Table 2, across environmental, social, and economic dimensions [22, 23]:

a) Environmental Benefits

Agroecological practices protect ecosystem resilience by increasing habitat diversity and reducing chemical inputs. They enhance soil health through crop rotation, cover crops, and organic amendments. Sustainable intensification methods improve soil structure, nutrient retention, and erosion [24]. Water conservation through drip irrigation, rainwater collection, and agroforestry reduces usage and improves water quality [25, 27]. Agroecology and sustainable intensification techniques improve climate resilience by encouraging varied agricultural systems, agroforestry, and soil carbon storage, reducing greenhouse fuel emissions and preparing for climate variability [25].

b) Social Benefits

Agroecological strategies enhance food security by increasing production and improving food diversity and nutritional quality. In addition with farmer-led research they promote farmer empowerment, cooperative decision-making, and statistics sharing [26]. Sustainable intensification aims for equity and social justice through minimizing differences in small farmers in their access to resources [28]. Agroecological structures also maintain cultural history by means of the use of conventional information and indigenous farming techniques, allowing promotion of community and cultural preservation [29-32].

c) Economic Benefits

Sustainable intensification methods increase farm productiveness through decreasing cost required for the inputs, increasing agricultural yield and quality, and maximizing resource use performance as factor of efficiency. Agroecology contributes to self-sufficiency and resilience in agricultural organizations via statistics sharing, farmer-led studies, and organization decision-making. Agro-ecological and environmentally friendly products frequently fetch premium expenses in specialised markets, incentivizing manufacturers to pursue sustainable practices [33-35]. These methods also increase the capacity of agricultural structures to cope with market uncertainty and climate-related challenges, exploiting different sources of income and reducing dependence on external inputs.

Table 2: Benefits of agroecology and sustainable intensification

Dimension	Benefit Type	Agroecology	Sustainable Intensification
Environmental	Conservation of Biodiversity	Increases habitat diversity and reduces chemical inputs	Similar benefits through diversified practices
	Soil Health	Enhances fertility with crop rotation, cover crops, organic amendments	Similar benefits through soil management strategies

	Water Conservation	Employs water-efficient strategies like drip irrigation, rainwater collection	Similar emphasis on water efficiency and quality improvement
	Climate Resilience	Improves resilience through diverse systems and carbon storage	Enhances resilience via diverse agricultural practices
Social	Food Security	Increases production, diversity, and nutritional quality	Similar benefits through intensified but sustainable practices
	Farmer Empowerment	Promotes self-sufficiency and resilience via cooperative decision-making	Similar empowerment but focuses more on technological and methodological innovations
	Fairness and Social Justice	Seeks inclusivity in agricultural development	Aims to minimize disparities in resource access
	Cultural Preservation	Maintains cultural heritage with traditional knowledge	May incorporate traditional practices but with a modern integration
Economic	Enhanced Farm Productivity	Reduces input costs and increases yields and quality	Increases productivity through optimized resource use
	Savings on Costs	Promotes resilience and self-sufficiency to reduce costs	Similar savings through efficiency and reduced reliance on inputs
	Market Opportunities	Commands premium prices in niche markets	Similar market opportunities for sustainable products
	Resilient Agricultural Systems	Increases resilience to market and climate challenges	Similar resilience through diversified strategies and reduced inputs dependency

Agroecological practices enhance atmosphere resilience by growing habitat diversity, lowering chemical inputs, and enhancing soil fitness. Sustainable intensification methods enhance soil quality, nutrient retention, erosion, and water conservation. Agroecological techniques additionally aid in food protection, put their focus on farmer empowerment, and save the cultural aspects. Sustainable agricultural strategies boom farm productiveness, reduce input expenses, and produces a growth in agricultural yield.

5. Conclusion

This comparative evaluation of agro-ecology and sustainable intensification focuses on the good contributions that both strategies can have on the potential to promote economic improvement, environmental sustainability, and international food protection. The use of sustainable intensification strategies, like integrated soil-crop system management and nitrogen management systems, has verified high-quality results in growing agricultural yields even as decreasing environmental degradation. On the other side, agro ecology will increase climate resilience along with providing a stable foundation for advancing social equity, economic viability, soil health, and biodiversity. The interactions between sustainable intensification and agroecology highlights the need for comprehensive and integrated strategies that use ecological ideas, technical advances and indigenous facts to create principles, abundant and sustainable food structures. Making sure environmental preservation, food protection, and socioeconomic equity can be carried out through means of combining the principles of agroecological and sustainable intensification. These desires are critical considering the continuous problems in feeding a growing global population as the challenge persists in the form of weather changes and finite natural resources. Comparative Analysis of Agro-Ecology and Sustainable Intensification has the following key outcomes:

- a) It can be concluded that Agro-ecology has the potential to enhance economic viability, soil health along with safeguarding biodiversity.
- b) The benefits of integration of the two strategies of agroecological and sustainable intensification includes environmental protection, security of food, and socioeconomic integrity.
- c) when looking at each method individually it can be concluded that each one of them can contribute to financial growth taking care of environmental sustainability, and global food security. The methods of sustainable intensification measures, integrated soil-crop system management and nitrogen control, has the ability to boost the agricultural output while lessening environmental impacts.

References

[1] Bernard, B., & Lux, A. “How to feed the world sustainably: an overview of the discourse on agroecology and sustainable intensification”. *Regional Environmental Change*, 17(5). (2017): 1279-1290.

[2] Dumont, B., Groot, J. C., & Tichit, M. “Make ruminants green again–how can sustainable intensification and agroecology converge for a better future.” *Animal*, 12(s2). (2018):s210-s219.

- [3] Mockshell, J., & Kamanda, J. "Beyond the agroecological and sustainable agricultural intensification debate: Is blended sustainability the way forward" *International Journal of Agricultural Sustainability*, 16(2). (2018):127-149.
- [4] Mishra, Pradumn, Urja Jain, Siddharth Choudhury, Surjeet Singh, Anish Pandey, Abhishek Sharma, Ramanpreet Singh, Vimal Kumar Pathak, Kuldeep K. Saxena, and Anita Gehlot. "Footstep planning of humanoid robot in ROS environment using Generative Adversarial Networks (GANs) deep learning." *Robotics and Autonomous Systems* 158 (2022): 104269.
- [5] Boincean, B., Dent, D., Boincean, B., & Dent, D. "Agroecology: Science for sustainable intensification of agriculture." *Farming the Black Earth: Sustainable and Climate-Smart Management of Chernozem Soils*, 19-27.
- [6] González de Molina, M., & Guzmán Casado, G. I. "Agroecology and ecological intensification. A discussion from a metabolic point of view". *Sustainability*, 9(1). (2017):86.
- [7] Wezel, A., Soboksa, G., McClelland, S., Delespesse, F., & Boissau, A.. "The blurred boundaries of ecological, sustainable, and agroecological intensification: a review". *Agronomy for sustainable development*, 35. (2015): 1283-1295.
- [8] Singh, P., Singh, A. K., Arun, V., & Tripathi, D. K. (2024). Comparative study of all-optical INVERTER and BUFFER gates using MZI structure. *Journal of Optical Communications*, 44(s1), s253-s261.
- [9] Alkorbi, A. S., Kumar, K. Y., Prashanth, M. K., Parashuram, L., Abate, A., Alharti, F. A., ... & Raghu, M. S. (2022). Samarium vanadate affixed sulfur selfdoped g-C3N4 heterojunction; photocatalytic, photoelectrocatalytic hydrogen evolution and dye degradation. *International Journal of Hydrogen Energy*, 47(26), 12988-13003.
- [10] Altieri MA, Nicholls CI, Montalba R. "Technological Approaches to Sustainable Agriculture at a Crossroads: An Agroecological Perspective". *Sustainability*. 9(3). (2017): 349.
- [11] Gangwar, D.S., Tyagi, S. & Soni, S.K. "A conceptual framework of agroecological resource management system for climate-smart agriculture". *Int. J. Environ. Sci. Technol.* 16. (2019): 4123–4132.
- [12] Srinivasan, K., Porkumaran, K., & Sainarayanan, G. (2009, August). Improved background subtraction techniques for security in video applications. In *2009 3rd International Conference on Anti-counterfeiting, Security, and Identification in Communication* (pp. 114-117). IEEE.
- [13] Vishwanatha, H. M., Kuldeep K. Saxena, Alokesh Pramanik, and Ajit Behera. "Cryo treatment and corrosion studies of nickel-titanium shape-memory alloy." *Proceedings of the Institution of Mechanical Engineers, Part E: Journal of Process Mechanical Engineering* (2023): 09544089231159250.
- [14] Leippert, F., Darmaun, M., Bernoux, M., Mpheshea, M., Müller, A., Geck, M., ... & Termote, C. "The potential of agroecology to build climate-resilient livelihoods and food systems" (2020).
- [15] Kumar, C. P., Raghu, M. S., Prathibha, B. S., Prashanth, M. K., Kanthimathi, G., Kumar, K. Y., ... & Alharthi, F. A. (2021). Discovery of a novel series of substituted quinolines acting as anticancer agents and selective EGFR blocker: Molecular docking study. *Bioorganic & Medicinal Chemistry Letters*, 44, 128118.
- [16] Kerr, R. B., Young, S. L., Young, C., Santoso, M. V., Magalasi, M., Entz, M., ... & Snapp, S. S. "Farming for change: developing a participatory curriculum on agroecology, nutrition, climate change and social equity in Malawi and Tanzania". In *Critical Adult Education in Food Movements*. (2022): 29-46. Cham: Springer Nature Switzerland.
- [17] Upadhyay, K. K., Srivastava, S., Mishra, N. K., & Shukla, N. K. (2020). Design and performance analysis of MZI based 2×2 reversible XNOR logic gate. *Journal of Optical Communications*, 41(3), 235-239.
- [18] Zaremba H, Elias M, Rietveld A, Bergamini N. "Toward a Feminist Agroecology". *Sustainability*. 13(20). (2021): 11244.
- [19] Buddi, Tanya, Susmitha Valli Gogula, and Kuldeep K. Saxena. "Manufacturing and Evaluation of Mechanical Properties for Rice Husk ParticleBoard Using IoT." *Indian Journal of Engineering and Materials Sciences (IJEMS)* 29, 6 (2023): 750-754.
- [20] Cassman, K. G., & Grassini, P. "A global perspective on sustainable intensification research". *Nature Sustainability*, 3(4). (2020): 262-268.
- [21] Burney, J. A., Davis, S. J., & Lobell, D. B. "Greenhouse gas mitigation by agricultural intensification". *Proceedings of the national Academy of Sciences*, 107(26). (2010):12052-12057.
- [22] Darjee, S., Shrivastava, M., Langyan, S., Singh, G., Pandey, R., Sharma, A., ... & Singh, R. "Integrated nutrient management reduced the nutrient losses and increased crop yield in irrigated wheat." *Archives of Agronomy and Soil Science* 69.8 (2023): 1298-1309.
- [23] Xia, L., Cao, L., Yang, Y., Ti, C., Liu, Y., Smith, P., ... & Yan, X. "Integrated biochar solutions can achieve carbon-neutral staple crop production." *Nature Food*, 4 (3) (2023): 236-246.
- [24] Raji, Atchudan, Jebakumar Immanuel Edison Thomas Nesakumar, Shanmugam Mani, Suguna Perumal, Vinodh Rajangam, Somanathan Thirunavukkarasu, and Yong Rok Lee. "Biowaste-originated heteroatom-doped porous carbonaceous material for electrochemical energy storage application." *Journal of Industrial and Engineering Chemistry* 98 (2021): 308-317.
- [25] Ashwini, S., Prashantha, S. C., Naik, R., & Nagabhushana, H. (2019). Enhancement of luminescence intensity and spectroscopic analysis of Eu³⁺ activated and Li⁺ charge-compensated Bi₂O₃ nanophosphors for solid-state lighting. *Journal of Rare Earths*, 37(4), 356-364.

- [26] Khan, Amir, and Kuldeep Kumar Saxena. "A review on enhancement of mechanical properties of fiber reinforcement polymer composite under different loading rates." *Materials Today: Proceedings* 56 (2022): 2316-2322.
- [27] Manohar, T., Prashantha, S. C., Nagaswarupa, H. P., Naik, R., Nagabhushana, H., Anantharaju, K. S., ... & Premkumar, H. B. (2017). White light emitting lanthanum aluminate nanophosphor: near ultra-violet excited photoluminescence and photometric characteristics. *Journal of Luminescence*, 190, 279-288.
- [28] Telagam, Nagarjuna, Nehru Kandasamy, and Menakadevi Nanjundan. "Smart sensor network based high quality air pollution monitoring system using labview." *International Journal of Online Engineering (iJOE)* 13,08 (2017): 79-87.
- [29] Numan, Arshid, Atal AS Gill, Saqib Rafique, Manisha Guduri, Yiqiang Zhan, Balaji Maddiboyina, Lijie Li, Sima Singh, and Nam Nguyen Dang. "Rationally engineered nanosensors: A novel strategy for the detection of heavy metal ions in the environment." *Journal of Hazardous Materials* 409 (2021): 124493.
- [30] Bhojak, Vishal, Jayahari Lade, Jinesh Kumar Jain, Amar Patnaik, and Kuldeep Kumar Saxena. "Investigation of annealing on CR-2 grade steel using Taguchi and Taguchi based gray relational analysis." *Advances in Materials and Processing Technologies* 8, (2022): 2231-2246.
- [31] Naresh, Muthunoori, and P. Munaswamy. "Smart agriculture system using IoT technology." *International journal of recent technology and engineering* 7, 5 (2019): 98-102.
- [32] Spandana, Kancheti, and VR Seshagiri Rao. "Internet of things (Iot) based smart water quality monitoring system." *International Journal of Engineering & Technology* 7, 3.6 (2018): 259-262.
- [33] Sankhla, Arvind M., Kaushik M. Patel, Mayur A. Makhesana, Kuldeep K. Saxena, and Nakul Gupta. "Experimental investigation of tool wear in machining of SiC based Al-MMC." *Advances in Materials and Processing Technologies* 8, (2022): 635-654.
- [34] Radhakrishna, Vangipuram, P. V. Kumar, V. Janaki, and N. Rajasekhar. "Estimating prevalence bounds of temporal association patterns to discover temporally similar patterns." In *International Conference on Soft Computing-MENDEL*, 209-220. Cham: Springer International Publishing, 2016.
- [35] Swarna, K. S. V., Vinayagam, A., Ananth, M. B. J., Kumar, P. V., Veerasamy, V., & Radhakrishnan, P. (2022). A KNN based random subspace ensemble classifier for detection and discrimination of high impedance fault in PV integrated power network. *Measurement*, 187, 110333.