A Survey on Signal processing tools for the analysis of heart rate variation: an approach for the early diagnosis and management of cardiovascular disease

Mangoné FALL[1*], *Bathie POUYE*[1], Ibrahima *GUEYE*[2] And *Idy DIOP*[1]

[1]Cheikh Anta Diop University (UCAD), Polytechnic Institute (ESP), Department of computer sciences, Dakar, Senegal
[2]Gaston Berger University (UGB), Department of Applied Physics, Saint Louis, Senegal

Abstract:

Heart rate variability (HRV) is a non-invasive, reliable, and reproducible biomarker of autonomic nervous system (ANS) function. This review provides a critical and comparative analysis of the primary HRV evaluation methods (temporal, frequency, non-linear, and geometric) as applied to the early diagnosis and management of cardiovascular, metabolic, psychiatric, and neurological pathologies. We place particular emphasis on evaluating the accuracy, robustness, and clinical applicability of these methods. Recent advances in signal processing and artificial intelligence (AI) are paving the way for more precise real-time detection tools using portable devices suitable for telemedicine. However, challenges such as protocol heterogeneity, the confounding effect of heart rate, data imbalance in AI models, and a lack of standardization still limit large-scale clinical deployment. This study not only highlights the promising prospects of integrating HRV into personalized prevention strategies but also provides a critical discussion of methodological challenges and suggests avenues for future improvement, including the integration of explainable AI and robust validation frameworks.

Key words: Heart rate variability (HRV), autonomic nervous system (ANS), signal processing, temporal analysis, frequency analysis, non-linear methods, telemedicine, cardiac monitoring.

1-Introduction:

Heart Rate Variability (HRV), defined as the variation in time intervals between consecutive heartbeats (RR intervals), has evolved from a simple research curiosity to a critical non-invasive tool for assessing autonomic nervous system (ANS) function [1].The dynamic interplay between the sympathetic and parasympathetic branches of the ANS imprints complex patterns on heart rhythm, which HRV quantifies.
The clinical relevance of HRV is well-established, with studies linking reduced HRV to various conditions, including coronary artery disease [2], hypertension, myocardial

Corresponding author: mangone.fall@esp.sn

infarction[3], and diabetic neuropathy[4]. Furthermore, its utility has expanded into psychiatry , geriatrics, and behavioral medicine.
Recent technological advances have enabled more accessible HRV monitoring through portable devices and modern acquisition systems such as photoplethysmography imaging (PPGI) and Gyrocardiography (GCG). These innovations support the growing integration of HRV into telemedicine and personalized healthcare platforms.

Despite its potential, the widespread clinical adoption of HRV faces significant hurdles. The field features a plethora of analysis methods, ranging from simple statistical measures in the time domain to complex non-linear and AI-based approaches, each with its own strengths, weaknesses, and interpretive caveats. This diversity, while a sign of a vibrant field, leads to challenges in reproducibility, comparability between studies, and clinical interpretation. Often, reviews tend to be descriptive, listing methods without providing a critical framework for their selection and application in specific contexts.
The primary objective of this review is therefore to move beyond a descriptive listing and provide a systematic, critical, and comparative analysis of existing HRV analysis methods. We aim to systematically present and critique major HRV analysis methods while evaluating their performance across multiple dimensions, including accuracy, robustness, computational complexity, and clinical applicability. We will specifically address methodological challenges in AI-based approaches, such as data imbalance and interpretability issues, and provide a practical framework through comparative tables and an integrative figure to guide method selection. Finally, we will outline future directions for standardizing HRV analysis and integrating it into telemedicine and personalized healthcare platforms.
Having established the clinical potential of HRV and the current methodological challenges, the following section presents the essential physiological foundations underlying the autonomic mechanisms that HRV aims to measure.

2. Physiological foundations and conceptual bases of VFC:

2.1. Definition and Physiological Mechanisms:

HRV is a biomarker of the autonomic nervous system's influence on the sinus node. The ANS, governed by a central autonomic network[5], continuously modulates heart rate through its sympathetic (acceleratory) and parasympathetic (decelerator) arms. Key mechanisms influencing HRV include:
Respiratory Sinus Arrhythmia (RSA): The parasympathetically mediated heart rate fluctuation linked to the respiratory cycle[6].
- Baroreflex Sensitivity: This critical feedback mechanism, involved in blood pressure regulation, continuously adjusts heart rate in response to vascular stretch, contributing primarily to the low-frequency components of HRV.
- Thermoregulation and Hormonal Influences: Slower regulatory processes reflected in the very low-frequency components of the HRV spectrum.



Fig.1: RR interval space [3]

As illustrated in Figure 1, the representation of RR intervals provides the fundamental input for quantifying these autonomic oscillations. A crucial methodological consideration is the mathematical confounding effect of the basal heart rate itself on HRV measures. As demonstrated by Billman et al [7], higher heart rates can mechanically reduce HRV indices independent of autonomic tone, necessitating the application of appropriate correction methods for accurate physiological interpretation [8].

2.2. Key HRV Parameters: A Synopsis:

HRV analysis methodologies can be broadly categorized into three primary domains, each offering unique insights into autonomic function:

- Time-Domain Analysis: This approach utilizes simple statistical measures derived directly from RR intervals (e.g., SDNN, RMSSD, pNN50) to quantify the overall magnitude of heart rate fluctuations over time.
- Frequency-Domain Analysis: Spectral methods (e.g., FFT, AR modeling, Wavelet Transform) decompose the HRV signal into its constituent oscillatory components. This allows for the examination of power within specific frequency bands (VLF, LF, HF), which are associated with different physiological regulatory processes.
- Non-Linear Domain Analysis: Derived from chaos and complexity theory, these methods (e.g., Entropy measures, Detrended Fluctuation Analysis, Poincare plots) aim to quantify the complexity and fractal-like properties of heart rate dynamics. They are often particularly sensitive to subtle, early pathological changes that may not be evident in linear analyses [9] [10].

This foundation in the origins and measurement domains of HRV sets the stage for a critical examination of the specific analytical methods, their comparative strengths, and their optimal clinical applications, which is the focus of the following section.

3. A Critical Comparison of HRV Analysis Methods:

This section provides a critical evaluation of the main HRV analysis methodologies, progressing from traditional to advanced approaches. Each method is assessed based on its fundamental principles, practical advantages, limitations, and optimal application contexts.

3.1. Time-Domain Analysis:

Principle & Indicators: Time-domain analysis involves the direct statistical or geometric examination of RR intervals. Common statistical measures include SDNN (reflecting overall

variability), RMSSD (representing parasympathetic activity), and pNN50 (quantifying the proportion of adjacent intervals differing by more than 50 ms). Geometric methods analyze the shape of RR interval histograms.
Advantages: The primary strengths of time-domain analysis are its computational simplicity, intuitive interpretation, and proven robustness for long-term monitoring (e.g., 24-hour recordings).
Limitations & Challenges: A significant limitation is its inability to distinguish between sympathetic and parasympathetic nervous activities. These measures are highly sensitive to artifacts and outliers. Furthermore, their reliability depends strongly on recording duration; for instance, SDNN is not recommended for short-term recordings.
Optimal Context: This approach is best suited for initial screening, long-term risk stratification (such as 24-hour SDNN assessment post-myocardial infarction), and fitness tracking applications where computational resources are limited.

3.2. Frequency-Domain Analysis:

Principle & Indicators: Frequency-domain methods decompose the HRV signal into its constituent frequency components using various spectral estimation techniques. The key bands of physiological interest are LF (0.04-0.15 Hz), often associated with sympathetic modulation and baroreflex activity, and HF (0.15-0.4 Hz), primarily influenced by parasympathetic (vagal) activity linked to respiration.



Fig. 2: Illustration of the frequency bands in spectral analysis [11] .

As shown in Figure 2, these frequency bands provide a visual representation of autonomic oscillations. The analysis pipeline for spectral HRV assessment is systematic: beginning with the raw ECG signal acquisition, followed by crucial preprocessing steps including digital filtering and signal standardization. Frequency analysis, typically employing Fast Fourier Transform (FFT) or Autoregressive (AR) modeling, is then performed to generate the power spectrum, from which relevant spectral components are identified for clinical interpretation.
Advantages: Spectral analysis allows for the potential distinction between sympathetic and parasympathetic influences under controlled conditions. AR modeling typically provides smoother spectra and better frequency resolution for short data segments compared to the classical FFT approach.
Limitations & Challenges: The fundamental assumption of stationarity represents a major limitation, making standard FFT unsuitable for non-stationary signals commonly encountered in real-life settings. The physiological interpretation of the LF band remains

controversial and should not be considered a pure sympathetic indicator. While Wavelet Transform offers powerful time-frequency analysis capabilities for non-stationary signals, it introduces complexity in selecting the appropriate mother wavelet and carries higher computational costs.

Optimal Context: FFT and AR methods are optimal for controlled, short-term resting assessments (e.g., tilt-table tests). Wavelet Transform is particularly valuable for analyzing transient events or dynamic conditions such as sleep apnea or stress induction protocols.



Fig. 3: Neuro-vegetative balance [12] .



Fig.4: ECG Signal Frequency Processing Diagram.

The diagram illustrates the complete process of ECG signal frequency processing, structured in four main steps. The first step involves the input signal, which is the raw ECG signal captured directly from the electrodes. This signal contains all the electrical information related to cardiac activity. Next comes preprocessing, which involves two essential operations. First, digital filtering is applied to remove unwanted noise and artifacts. On the other hand, a normalization is carried out in order to homogenize the amplitude of the signal, thus guaranteeing a better quality of the data for the following analysis. The third step is frequency analysis, in which the Fast Fourier Transform (FFT) is used to decompose the signal into its different frequency components. Examining the spectrum allows us to identify the characteristic frequencies of the cardiac signal. Finally, feature extraction allows the collection of relevant information for medical analysis. Each step is linked to the next by a logical flow from the initial recording of the signal to obtaining the parameters useful for clinical interpretation. The Fast Fourier Transform (FFT) allows the spectral decomposition of the VFC signal, offering the advantage of rapid identification of frequency bands. However, it has limitations, including the need for a stationary signal and a lack of temporal information. In contrast, the wavelet transform (WT) is a time-frequency approach suitable for non-stationary signals, allowing for better analysis of transient variations in HRV. However, this method is more computationally complex. Wavelets were born from a need to analyze signals locally. Their origins date back to several scientific works, but one of the most notable was that of Jean Morlet in 1975, who, in the field of geophysics, proposed an approach inspired by Gabor. Unlike Fourier, Morlet varied the window size (instead of the number of oscillations), which allowed for better frequency and time resolution. Continuous

wavelet transforms (CWT): TOC is based on the use of a function called the mother wavelet, dilated and translated through parameters (a: scale, b: translation). It allows multiresolution analysis of the signal. This method respects admissibility conditions guaranteeing lossless reconstruction. TOC measures the similarity between the signal and distorted versions of the mother wavelet, which is valuable for analyzing signals with variable frequency content. For practical application, we use the discrete wavelet transform, where the parameters a and b are discretized. We adopt dyadic (logarithmic) sampling which avoids redundancies while preserving essential information. The DWT allows an efficient representation of the signal and facilitates reconstruction, with a good compromise between complexity and fidelity. Wavelets are currently used in many fields: image compression, signal denoising, fractal analysis, etc. In filtering, conjugate mirror filters (high-pass and low pass) are used to decompose and reconstruct the signal at different resolution levels. These filters are essential for perfect reconstruction, without data loss. Wavelet filtering offers a powerful solution suitable for non-stationary signals. It outperforms the Fourier transform in many situations, particularly for biomedical signals, thanks to its ability to provide a multi-resolution time-frequency representation. It is now an essential tool in modern signal processing.

3.3. Non-Linear and Fractal Analysis:

Principle & Indicators: Non-linear methods quantify the complexity and self-similarity (fractal properties) of HRV signals. Key techniques include Detrended Fluctuation Analysis (DFA), which quantifies long-range correlations; Sample Entropy (SampEn), measuring signal regularity and predictability; and Poincare plots (with indices SD1, SD2), which visualize and quantify beat-to-beat variability.

Advantages: These methods are highly sensitive to early pathological states that reduce system complexity, such as congestive heart failure and diabetes. They capture aspects of autonomic regulation that are typically invisible to conventional linear methods.

Limitations & Challenges: A significant barrier to clinical adoption is the lack of standardized parameters for calculations (particularly for entropy measures) and the difficult physiological interpretation of the resulting indices. The results can be highly sensitive to data length and noise, requiring significant expertise for proper implementation and interpretation.

Optimal Context: Non-linear analysis is most appropriately used in research settings for early disease detection, risk stratification in chronic conditions, and investigating the fundamental complexity of cardiac control mechanisms.

Table 1. Classical Methods :

Types of methods	Subcategory / Example	Description	Key Indicator	Typical recording time
Statistics (Time)	Averages, deviations	Analysis of RR intervals (CI)	SDNN, RMSSD, NN50, pNN50, CVr	5 min to 24 h
Geometric	Histograms, triangular index, scattergram	CI distribution	Mode, mode amplitude, index triangular, TINN, Scattergram area	5 min
Spectral (Frequenti She)	Parametric ue (Auto regression) - Non Parametric ue (FFT, Welch, Lomb)	Frequency component analysis	HF (0.15 0.4 Hz), LF (0.04–0.15 Hz), VLF (0.003–0.04 Hz), ULF (	5 min to 24 h

Table 2. **Non-Linear Methods Table:**

Methods	Description	Key indicators	Recommended duration
Hurst analysis (RS Analysis)	Measure, the relationship between trend and noise (fractality)	Hurst exponent (H)	≥ 5 min
Fluctuation analysis (DFA)	Analysis of self-similarity of time series	Fluctuation coefficient (α)	5 min to 24 h
Deceleration capacity / Acceleration	Anchoring of heart rate change points heart rate	Deceleration and acceleration coefficients	24 h
Wavelet transform	Localised time-frequency analysis, adapted to non-stationary signals	Wavelet spectral power (Daubechies)	Variable

Table 3. Methods for Heart Rate Variability (HRV) Analysis Key-Studies, Contribution and Limitations

Authors	Methods / Techniques	Contributions	Limits
Akselrod (1981) and al[13]	Spectral analysis (LF/HF)	First use of spectral analysis to assess autonomic activity via VFC	Based on assumptions of linearity, stationarity and Gaussian distribution
Pomeranz and al. (1985) [14]	Advanced spectral analysis	Evaluation of the bands LF/HF for sympathetic and parasympathetic regulation	Lack of sensitivity to detect non-linear dynamic complexity
Costa et al (2005, 2008)[15]	Multiscale Entropy (MSE), Approximate Entropy (ApEn)	Evaluation of VFC signal complexity on different time scales	Sensitive to the choice of parameters; loss of information in the long-term information loss

Ivanov et al (1999) [16],	Fractal and multifractal analysis	Demonstration of the multifractality of human VFC signals	Strong hypothesis of free-scale dynamics; difficult to validate clinically
Kiyono and al (2006, 2008) [17]	Analysis of non-Gaussianity	Non-Gaussianity indices for cardiac risk stratification	Based on log-normal assumption; sensitive to artefacts
Wendt and al (2018) [18]	Multiscale non-Gaussian representations with p-leaders	Robust and flexible approach, high discriminative capacity (CHF: survivors vs. nonsurvivors)	Mathematical complexity; requires broader clinical validation
Gallardo and al (2022)[19]	Geometric analysis using the Poincare representation (SD1, SD2, SD21)	SD21 used as LF/HF analogue. Good correlation with 5 min indices from 120 s of measurement.	Requires careful pre-processing, sensitivity to artefacts, possible overestimation of indices.
Nardelli and al. (2018)[20]	Analysis of SD12 (inverse of SD21) in simulated series from 15 to 60 s.	Very high correlation (ρ=0.994) for series $\geq$ 35 s with 1 h series.	On real data (27 subjects), accuracy decreases (ρ=0.961; e=14.47%). Validity limited to certain contexts.
Pecchia and al. (2018) [21]	Assessment critical of VFC-UC methods	Review of existing methods; recommendation of a rigorous approach to Concurrent validation approach	Underlines the absence of clear standards for VFC-UC. Approach still theoretical.
Malik and al (1996)[22]	International standards for VFC analysis (5 min and 24h)	Basic reference for standardised VFC assessment	Does not cover the ultra-short term; linear and stationary stationary methods only.
Hamed Azami and al [23]	SSA filter (Singular Spectrum Analysis)	Decomposition of EEG/ECG signals via SVD; Effective separation noise/signal separation; useful for non-stationary signals	Sensitivity to parameters (window length, number of components); risk of loss of information

Schmidt (NASA) (1960), Kalman [24]	Kalman filter	Recursive optimal filtering; applicable to dynamic linear systems; useful for prediction/monitoring	Requires an accurate model; less effective in the case of non-linearity; signal may be attenuated
Sameni R and al [25]	Extended Kalman Filter (EKF)	Adaptation for non-linear systems via linearisation (Taylor expansion)	Accuracy reduced by approximations; high algorithmic complexity
Wavelet Transform (DWT)	Wavelet Transform (DWT)	Non-stationary analysis; good time frequency Frequency localisation; preservation of signal shape	Sensitivity to choice of wavelet; risk of degradation at low SNR
Fourier (1768-1830)[26].	Fourier transform	Global frequency analysis; basis for analysis of deterministic signals	Not suitable for non-stationary signals; no local temporal information

Advanced methods and signal processing:

3.4. AI and Machine Learning-Based Analysis:

Principle & Techniques: This emerging approach utilizes machine learning algorithms (e.g., Support Vector Machines, Random Forests) and deep learning models (e.g., Convolutional Neural Networks, Long Short-Term Memory networks) to automatically identify complex, non-linear patterns in HRV data for classification or prediction tasks.

Advantages: Potential for superior accuracy in detecting specific pathologies by learning from high-dimensional data. Ability to integrate HRV with other clinical or sensor data.

Limitations & Methodological Challenges: Data Imbalance: Models often perform poorly on underrepresented classes (e.g., rare cardiac events).

Overfitting: High risk of models learning noise from the training data instead of generalizable patterns, especially with small datasets. Interpretability ("Black Box"): The lack of transparency in how deep learning models reach decisions is a major barrier to clinical trust and adoption. Techniques like SHAP or LIME are needed.

Generalization: Models trained on one dataset with specific acquisition protocols often fail on data from different sources.

Optimal Context: Large-scale data analysis, development of diagnostic support systems, and personalized risk prediction where large, well-annotated datasets are available.

Table 4. Historical Evolution and Key and Milestones in Heart Rate Variability (HRV) Research-Contributions and Limitations

Author/References	Years	Contributions	Limits
Ebers Papyrus[27]	1550 BC	First mention of heartbeats	No link established with the autonomic nervous system

Aristotle, Hippocrates[28]	Ancient Greece	Interest in the heart and circulation	Philosophical and descriptive approach
Wang Shu-He [29]	265-317	Link between pulses and pathologies	Empirical interpretation
Floyer[30]	1707	Invented a watch to measure pulse	Limited to manual observation
Hon, Lee, Wolf	1963-1975	VFC in gynaecology, foetal stress, sudden death	Limited to certain populations
Sayers, Kahneman, Porges	1970s	Use of VFC in psychology	Methods analytical still rudimentary
Akselrod and al[13]	1981	HF reflects the parasympathetic	Requires precise spectral measurements
Malliani and Pagani[31]	1994-1995	LF associated with sympathetic	Controversy over the exact meaning of the bands
Akselrod, Pomeranz, Koh[32], [14]	2000s	Refine interpretation of bands LF/HF	Need for clinical validation
Kurths et al [11]	1995	Non-linear methods for VFC	Complexity of methods
Carnethon and al[2]	2002	Link between VFC and neurovegetative imbalances	Study on healthy population only
Menezes and al[33]	2004	VFC reduced in hypertensive patients	No longitudinal analysis
Terathongkum and al [34]	2004	VFC predicts hypertension	Correlation but no causality established
Bittencourt and al [35]	2005	Parasympathetic increase with controlled breathing	Controlled experimental conditions
Pecyna [36]	2006	Decreased VFC post-infarction	Lack of mechanistic data
Carney and al [37]	2007	Link between inflammation and FVC	Correlational study
Karas and al [12]	2008	Decreased LF in elderly hypertensive subjects	Specific to elderly subjects
Larosa and al[38]	2008	Decreased VFC in myocardial infarction	Context specific to myocardial ischaemia
Task Force of the ESC & NASPE[3]	1996	Establishes standards for HRV analysis	Does not explicitly address uncertainties.

Peltola and al	2012	Shows the mismatch between short-term and long term	No solution for uncertainty related to segmentation.
Berntson and al [1]	1997	Recommends frequency ranges for spectral analysis	Not focused on modern technical uncertainties.
Clifford and al [39]	2006	Details ECG artefacts and suggests treatments	No formal modelling of uncertainties.
Mateo and Laguna[40]	2003	Effect of sampling frequency on HRV	Analysis limited to a few indices.
Clifford and al[13]	2006	Combines several QRS detectors for greater robustness	Requires a lot of computational resources
Schafer & Vagedes	2013	Investigate the effect of sampling on PRV	Does not generalise to all HRV.
Sassi and al[41]	2015	Minimum recommended durations for ultra-short HRV	Ultra-short remains unreliable for some indices.
Aboy and al [42]	2005	Compares interpolation methods for missing data	Simulated study, not under real conditions.
Schäfer and al[43]	1998	Spectro-temporal analysis for HRV	Less used clinically.
Al-Nashash and al [44]	2000	Kalman filtering for noisy HRV	Complex model to calibrate.
Lu and al[45]	2012	Impact of QRS errors on HRV (Monte Carlo)	Does not cover all uncertainties (e.g. physiological).
Zhang and al[46]	2021	HRV estimated by deep learning	Clinical validation still required.

This table illustrates the evolution of knowledge and investigation techniques of VFC over the centuries.

Table 5: Comparative Analysis of HRV Methods:

Method	Accuracy/Robustness	Computational complexity	Optimal Application Context	Key Limitations
Time-Domain	Moderate, sensitive to artifacts	Low	Long-term monitoring, fitness tracking, initial screening	Cannot dissect SNS/PNS activity; less detailed.

Frequential-Domain (FFT/AR)	High for stationary signals	Low to Moderate	| Assessment of sympathovagal balance (rest/tilt)	Assumes stationarity; controversial LF interpretation
Frequential-Domain (Wavelets)	High for non-stationary signals	High	Analysis of transient events (sleep, stress)	Complex parameter selection (mother wavelet)
Non-Linear	High sensitivity, but parameter-dependent	Moderate to High	Early disease detection, complexity assessment	Complex interpretation; lacks clinical standardization
AI/ML	Potentially very high, but context-dependent	Very High	Pattern recognition, diagnostic support, prediction	"Black-box" nature; requires large, annotated datasets; risk of overfitting.

4. Discussion: Synthesis, Challenges, and Future Directions:

Our comprehensive comparative analysis reveals a fundamental principle: there is no single "best" method for HRV analysis. The optimal choice represents a strategic trade-off based on the specific clinical question, signal quality constraints, available computational resources, and required interpretability. Time-domain methods offer straightforward accessibility for basic screening, frequency-domain analysis provides valuable physiological insights under controlled conditions, non-linear methods capture hidden complexity indicative of early pathology, while AI approaches promise powerful pattern recognition capabilities at the potential cost of transparency.

4.1. Overcoming Current Methodological Challenges:

Several critical challenges must be systematically addressed to facilitate the transition of HRV from research to widespread clinical practice:

Standardization: There is an urgent need for universally accepted protocols governing data acquisition, preprocessing pipelines, and parameter settings, particularly for non-linear and AI methodologies. This standardization is a fundamental prerequisite for meaningful multi-center studies and robust clinical translation.

Heart Rate Correction: The confounding effect of basal heart rate on HRV indices must be systematically addressed in analytical workflows. Correction techniques, such as those proposed by Billman [13], should become a standard procedural step to ensure accurate physiological interpretation.

AI-Specific Hurdles: Future development must prioritize explainable AI (XAI) models that provide transparent decision-making pathways. Concurrently, the creation of large, public, and meticulously curated HRV databases is essential, alongside the implementation of advanced techniques to rigorously address data imbalance and overfitting.



Fig.5: Comprehensive overview of HRV analysis methods and their clinical applications

The diagram illustrates the complete workflow from signal acquisition to clinical implementation, highlighting the four main analytical domains along with their respective strengths and optimal usage contexts. As summarized in this figure, the choice of HRV analysis methodology should be guided by the specific clinical question, available resources, and technical constraints, with each domain offering complementary insights into autonomic function.

4.2. Future Directions:

The evolving landscape of HRV analysis points toward several promising research directions:

Hybrid Models: Combining features from linear, non-linear, and frequency domains as input to machine learning classifiers represents a promising avenue for developing more robust and interpretable diagnostic tools.

Integration with Telemedicine: The development of low-power, computationally efficient algorithms for real-time HRV analysis on wearable devices (utilizing PPG, SCG, and GCG sensors) will be crucial for enabling continuous, remote monitoring in ambulatory settings.

Personalized Medicine: HRV trends, when combined with other digital biomarkers, can be integrated into dynamic personalized health dashboards. This approach enables dynamic risk assessment and preemptive intervention strategies in chronic disease management, moving toward truly personalized healthcare.

The successful addressing of these challenges and the pursuit of these future directions will determine the clinical impact of HRV analysis in the coming decade, potentially establishing it as a cornerstone of modern, data-driven cardiovascular care.

5. Conclusion:

Heart rate variability analysis stands at a pivotal juncture, having matured from a research tool into a potent, non-invasive biomarker with wide-ranging applications from cardiology to psychiatry. This review has provided a critical framework for understanding the landscape of HRV methodologies, highlighting that its future clinical utility hinges not on the discovery of new indices, but on the critical selection, rigorous validation, and intelligent integration of existing ones. The analysis demonstrates that HRV represents a mature non-invasive biomarker whose clinical utility extends well beyond cardiology. Future impact depends on critical selection and rigorous validation of existing methods rather than development of new indices. Large-scale deployment requires overcoming standardization challenges, basal heart rate correction, and developing transparent, robust AI systems. By tackling these challenges of standardization, methodological rigor, and transparent AI, HRV can fully transition into a cornerstone of modern, data-driven, and personalized healthcare, particularly through integration with telemedicine platforms and personalized medicine approaches.

References:

[1] G. G. Berntson et al., "Heart rate variability: Origins, methods, and interpretive caveats," *Psychophysiology*, vol. 34, no. 6, pp. 623-648, 1997, doi: 10.1111/j.1469-8986. 1997.tb02140. x.

[2] M. R. Carnethon et al., "Does the Cardiac Autonomic Response to Postural Change Predict Incident Coronary Heart Disease and Mortality: The Atherosclerosis Risk in Communities Study," *American Journal of Epidemiology*, vol. 155, no. 1, pp. 48-56, Jan. 2002, doi: 10.1093/aje/155.1.48.

[3] T. Wu, V. Blazek, and H. J. Schmitt, "Photoplethysmography imaging: a new noninvasive and noncontact method for mapping of the dermal perfusion changes," in *Optical Techniques and Instrumentation for the Measurement of Blood Composition, Structure, and Dynamics*, SPIE, 2000, pp. 62-70. Accessed: Jun. 16, 2025. [Online]. Available: https://www.spiedigitallibrary.org/conference-proceedings-of-spie/4163/1/Photoplethysmography-imaging--a-new-noninvasive-and-noncontact-method-for/10.1117/12.407646.short

[4] M. G. Morshed, M. A. Mukit, K. I. U. Ahmed, R. Mostafa, S. Parveen, and A. H. Khandoker, "Heart rate variability analysis for diagnosis of diabetic peripheral neuropathy," in *2020 IEEE Region 10 Symposium (TENSYMP)*, Dhaka, Bangladesh: IEEE, 2020, pp. 1253-1256. doi: 10.1109/TENSYMP50017.2020.9230736.

[7] A. S. Juarascio, R. J. Crochiere, T. M. Tapera, M. Palermo, and F. Zhang, "Momentary changes in heart rate variability can detect risk for emotional eating episodes," *Appetite*, vol. 152, p. 104698, 2020, Accessed: Jun. 16, 2025. [Online]. Available: https://www.sciencedirect.com/science/article/pii/S0195666319311808

[8] W. Verkruysse, L. O. Svaasand, and J. S. Nelson, "Remote plethysmographic imaging using ambient light," *Optics Express*, vol. 16, no. 26, pp. 21434-21445, 2008, Accessed: Jun. 16, 2025. [Online]. Available: https://opg.optica.org/abstract.cfm?uri=oe-16-26-21434

[9] F. Shaffer and J. P. Ginsberg, "An overview of heart rate variability metrics and norms," *Frontiers in Public Health*, vol. 5, p. 258, 2017, Accessed: Jun. 16, 2025. [Online]. Available: https://www.frontiersin.org/articles/10.3389/fpubh.2017.00258/full

[5] E. E. Benarroch, "The central autonomic network: functional organization, dysfunction, and perspective," in *Mayo Clinic Proceedings*, Elsevier, 1993, pp. 988-1001. Accessed: Jul. 15, 2025. [Online]. Available: https://www.sciencedirect.com/science/article/pii/S0025619612622721

[6] D. Servant, R. Logier, Y. Mouster, and M. Goudemand, "Heart rate variability: Interest in psychiatry," *L'Encéphale*, vol. 35, no. 5, pp. 423-428, Oct. 2009, doi: 10.1016/j.encep.2008.06.016.
[7] G. E. Billman, "The effect of heart rate on the heart rate variability response to autonomic interventions," *Frontiers in Physiology*, vol. 4, p. 222, 2013, Accessed: Jun. 16, 2025. [Online]. Available: https://www.frontiersin.org/articles/10.3389/fphys.2013.00222/full
[8] N. Lunardon, G. Menardi, and N. Torelli, "ROSE: a package for binary imbalanced learning," 2014, Accessed: Jun. 16, 2025. [Online]. Available: https://digitalcommons.unl.edu/r-journal/418/
[9] M. W. Peng, "Firm Growth in Transitional Economies: Three Longitudinal Cases from China, 1989-96," *Organization Studies*, vol. 18, no. 3, pp. 385-413, May 1997, doi: 10.1177/017084069701800303.
[10] T. Hotta, "Orbital ordering phenomena in d- and f-electron systems," *Reports on Progress in Physics*, vol. 69, no. 7, p. 2061, 2006, Accessed: Jun. 17, 2025. [Online]. Available: https://iopscience.iop.org/article/10.1088/0034-4885/69/7/R02/meta
[11] J. Cassirame, "Interest and limitations of heart rate variability analysis for sports practice," Ph.D. dissertation, Université de Franche-Comté, 2015. Accessed: Jun. 14, 2025. [Online]. Available: https://theses.hal.science/tel-01299801
[12] J. Marsac, "Heart rate variability: a cardiometabolic risk marker in public health," *Bulletin de l'Académie Nationale de Médecine*, vol. 197, no. 1, pp. 175-186, Jan. 2013, doi: 10.1016/S0001-4079(19)31635-8.
[13] S. Akselrod, D. Gordon, F. A. Ubel, D. C. Shannon, A. C. Berger, and R. J. Cohen, "Power Spectrum Analysis of Heart Rate Fluctuation: A Quantitative Probe of Beat-to-Beat Cardiovascular Control," *Science*, vol. 213, no. 4504, pp. 220-222, Jul. 1981, doi: 10.1126/science.6166045.
[14] B. Pomeranz et al., "Assessment of autonomic function in humans by heart rate spectral analysis," *American Journal of Physiology-Heart and Circulatory Physiology*, vol. 248, no. 1, pp. H151-H153, Jan. 1985, doi: 10.1152/ajpheart.1985.248.1.h151.
[15] M. S. R. Ferreira et al., "Surveillance of norovirus infections in the state of Rio De Janeiro, Brazil 2005–2008," *Journal of Medical Virology*, vol. 82, no. 8, pp. 1442-1448, Aug. 2010, doi: 10.1002/jmv.21831.
[16] P. C. Ivanov et al., "Sleep-wake differences in scaling behavior of the human heartbeat: analysis of terrestrial and long-term space flight data," *Europhysics Letters*, vol. 48, no. 5, p. 594, 1999, Accessed: Jul. 10, 2025. [Online]. Available: https://iopscience.iop.org/article/10.1209/epl/i1999-00525-0/meta
[17] K. Kiyono, J. Hayano, S. Kwak, E. Watanabe, and Y. Yamamoto, "Non-gaussianity of low frequency heart rate variability and sympathetic activation: lack of increases in multiple system atrophy and Parkinson disease," *Frontiers in Physiology*, vol. 3, p. 34, 2012, Accessed: Jul. 10, 2025. [Online]. Available: https://www.frontiersin.org/articles/10.3389/fphys.2012.00034/full
[18] M. B. Winn, D. Wendt, T. Koelewijn, and S. E. Kuchinsky, "Best Practices and Advice for Using Pupillometry to Measure Listening Effort: An Introduction for Those Who Want to Get Started," *Trends in Hearing*, vol. 22, Jan. 2018, doi: 10.1177/2331216518800869.
[19] D. Farge et al., "2022 international clinical practice guidelines for the treatment and prophylaxis of venous thromboembolism in patients with cancer, including patients with COVID-19," *The Lancet Oncology*, vol. 23, no. 7, pp. e334-e347, 2022, Accessed: Jul. 10, 2025. [Online]. Available: https://www.thelancet.com/journals/lanonc/article/PIIS1470-2045(22)00160-7/abstract
[20] A. Vitagliano et al., "Uterine fibroid size modifications during pregnancy and puerperium: evidence from the first systematic review of literature," *Arch Gynecol Obstet*, vol. 297, no. 4, pp. 823-835, Apr. 2018, doi: 10.1007/s00404-017-4621-4.

[21] L. Pecchia, R. Castaldo, L. Montesinos, and P. Melillo, "Are ultra-short heart rate variability features good surrogates of short-term ones? State-of-the-art review and recommendations," *Healthcare Technology Letters*, vol. 5, no. 3, pp. 94-100, Jun. 2018, doi: 10.1049/htl.2017.0090.
[22] M. Malik, "Heart Rate Variability: Standards of Measurement, Physiological Interpretation, and Clinical Use: Task Force of The European Society of Cardiology and the North American Society for Pacing and Electrophysiology," *Noninvasive Electrocardiology*, vol. 1, no. 2, pp. 151-181, Apr. 1996, doi: 10.1111/j.1542-474X.1996.tb00275.x.
[23] H. Azami and S. Sanei, "Spike detection approaches for noisy neuronal data: assessment and comparison," *Neurocomputing*, vol. 133, pp. 491-506, 2014, Accessed: Jul. 10, 2025. [Online]. Available: https://www.sciencedirect.com/science/article/pii/S0925231214000022
[24] M. S. Grewal and A. P. Andrews, "Applications of Kalman filtering in aerospace 1960 to the present [historical perspectives]," *IEEE Control Systems Magazine*, vol. 30, no. 3, pp. 69-78, 2010, Accessed: Jul. 10, 2025. [Online]. Available: https://ieeexplore.ieee.org/abstract/document/5466132/
[25] R. Sameni, M. B. Shamsollahi, and C. Jutten, "Filtering electrocardiogram signals using the extended Kalman filter," in *2005 IEEE Engineering in Medicine and Biology 27th Annual Conference*, IEEE, 2006, pp. 5639-5642. Accessed: Jul. 10, 2025. [Online]. Available: https://ieeexplore.ieee.org/abstract/document/1615765/
[26] J. Fourier, "When Joseph Fourier (1768-1830) and Erwin Schrödinger (1887-1961) are not on the same wavelength...", Accessed: Jul. 10, 2025. [Online]. Available: https://dial.uclouvain.be/pr/boreal/object/boreal%3A189166/datastream/PDF_01/view
[27] B. Ziskind, "Cardiovascular examination in light of the medical papyri of ancient Egypt," *Histoire des Sciences Médicales*, vol. 40, no. 1, p. 61, 2006, Accessed: Jul. 10, 2025. [Online]. Available: https://numerabilis.u-paris.fr/ressources/pdf/sfhm/hsm/HSMx2006x040x001/HSMx2006x040x001x0061.pdf
[28] J. Bidez and G. Leboucq, "An ancient anatomy of the human heart: Philistion of Locris and Plato's 'Timaeus'," *Revue des Études Grecques*, pp. 7-40, 1944, Accessed: Jul. 10, 2025. [Online]. Available: https://www.jstor.org/stable/44271994
[29] P. Zhang, T. Roberts, B. Richards, and L. J. Haseler, "Utilizing heart rate variability to predict ICU patient outcome in traumatic brain injury," *BMC Bioinformatics*, vol. 21, no. S17, Dec. 2020, doi: 10.1186/s12859-020-03814-w.
[30] P. Huard and M. Wong, "French investigations into Chinese science and technology in the 18th century," *Bulletin de l'École française d'Extrême-Orient*, vol. 53, no. 1, pp. 137-226, 1966, Accessed: Jul. 10, 2025. [Online]. Available: https://www.jstor.org/stable/43733267
[31] S. Ezzedini, "Study of temporal preparation: contribution of executive functions and the autonomic nervous system to motor and cognitive action control," Ph.D. dissertation, Université de Nanterre-Paris X, 2022. Accessed: Jul. 10, 2025. [Online]. Available: https://theses.hal.science/tel-04123669/
[32] A. J. R. van Gestel et al., "Predicting Daily Physical Activity in Patients with Chronic Obstructive Pulmonary Disease," *PLOS ONE*, vol. 7, no. 11, p. e48081, Nov. 2012, doi: 10.1371/journal.pone.0048081.
[33] L. Menezes Falcão, F. Pinto, L. Ravara, and P. A. van Zwieten, "BNP and ANP as diagnostic and predictive markers in heart failure with left ventricular systolic dysfunction," *J Renin Angiotensin Aldosterone Syst*, vol. 5, no. 3, pp. 121-129, Sep. 2004, doi: 10.3317/jraas.2004.028.
[34] S. Terathongkum and R. H. Pickler, "Relationships among heart rate variability, hypertension, and relaxation techniques," *Journal of Vascular Nursing*, vol. 22, no. 3, pp. 78-82, Sep. 2004, doi: 10.1016/j.jvn.2004.06.003.

[35] E. Hulten, S. Aslam, M. Osborne, S. Abbasi, M. S. Bittencourt, and R. Blankstein, "Cardiac sarcoidosis—state of the art review," *Cardiovascular Diagnosis and Therapy*, vol. 6, no. 1, pp. 50-63, Feb. 2016, doi: 10.3978/j.issn.2223-3652.2015.12.13.
[36] E. R. Bojorges-Valdez, J. C. Echeverria, R. Valdes-Cristerna, and M. A. Pena, "Scaling patterns of heart rate variability data," *Physiological Measurement*, vol. 28, no. 6, p. 721, 2007, Accessed: Jul. 10, 2025. [Online]. Available: https://iopscience.iop.org/article/10.1088/0967-3334/28/6/010/meta
[37] C. A. Stratakis, L. S. Kirschner, and J. A. Carney, "Clinical and molecular features of the Carney complex: diagnostic criteria and recommendations for patient evaluation," *The Journal of Clinical Endocrinology & Metabolism*, vol. 86, no. 9, pp. 4041-4046, 2001, Accessed: Jun. 16, 2025. [Online]. Available: https://academic.oup.com/jcem/article-abstract/86/9/4041/2848323
[38] C. Johnson et al., "Comparison of the efficacy of atorvastatin 10 mg versus 80 mg in reducing major cardiovascular events and repeated revascularizations in patients who have previously undergone percutaneous coronary intervention (post-hoc analysis of the TNT [Treating to New Targets] study)," *The American Journal of Cardiology*, vol. 102, no. 10, pp. 1312-1317, Nov. 2008, doi: 10.1016/j.amjcard.2008.07.023.
[39] B. Nassima and H. Alima, "R-wave detection in an electrocardiogram based on wavelet transform," Ph.D. dissertation, Université Mouloud Mammeri, 2012. Accessed: Jul. 10, 2025. [Online].
Available: https://www.ummto.dz/dspace/bitstream/ummto/8244/1/BouheraouaNassima_HammadiA.pdf
[40] J. Mateo and P. Laguna, "Improved heart rate variability signal analysis from the beat occurrence times according to the IPFM model," *IEEE Trans. Biomed. Eng.*, vol. 47, no. 8, pp. 985-996, Aug. 2000, doi: 10.1109/10.855925.
[41] R. Sassi et al., "Advances in heart rate variability signal analysis: joint position statement by the e-Cardiology ESC Working Group and the European Heart Rhythm Association co-endorsed by the Asia Pacific Heart Rhythm Society," *Europace*, vol. 17, no. 9, pp. 1341-1353, 2015, Accessed: Jul. 10, 2025. [Online]. Available: https://academic.oup.com/europace/article-abstract/17/9/1341/627516
[42] A. IDAOUI and K. Y. RISKOU, "Evaluation and adjustment of 'TRMM' precipitation products using artificial intelligence methods," Ph.D. dissertation, Université Ghardaia, 2018. Accessed: Jul. 10, 2025. [Online]. Available: http://dspace.univ-ghardaia.edu.dz/xmlui/handle/123456789/5033
[43] K. Schäfer, "Accelerated solvent extraction of lipids for determining the fatty acid composition of biological material," *Analytica Chimica Acta*, vol. 358, no. 1, pp. 69-77, 1998, Accessed: Oct. 18, 2025. [Online]. Available: https://www.sciencedirect.com/science/article/pii/S0003267097005874
[44] H. Al-Nashash, "Cardiac arrhythmia classification using neural networks," *Technology and Health Care*, vol. 8, no. 6, pp. 363-372, Dec. 2000, doi: 10.3233/thc-2000-8605.
[45] A. Giroud, "Validation study of heart rate and heart rate variability measurement devices," 2023, Accessed: Jul. 10, 2025. [Online]. Available: https://folia.unifr.ch/documents/326521/files/TM_Giroud_Alexandre.pdf
[46] C. S. L. Borrego, "Machine learning for the prediction of infection and evaluation of maturation in premature infants combining cardiac and respiratory variability," Ph.D. dissertation, Université de Rennes, 2021. Accessed: Jul. 10, 2025. [Online]. Available: https://theses.hal.science/tel-03365819/